

Due 3/27/2026, 9:30 A.M.

Solve the following problems and staple your solutions to this cover sheet. (Computer outputs must be put in the appropriate place in the solution, not attached as an appendix. You may physically cut and paste the output in the problem or allow appropriate space in the printout to add your hand written work.)

1. Sec 7.4 #3

2. Sec 7.4 #25

3. Sec 7.5 #4

4. Sec 7.5 #10

5. Sec 7.5 #29

6. Sec 7.6 #8

Hint: $\mathcal{L}\{u(t-a)g(t)\} = e^{-as}\mathcal{L}\{g(t+a)\}$.

7. Sec 7.6 #11

8. Sec 7.6 #15

9. Sec 7.6 #29

Hint: You may graph the solution using Mathematica.

10. Sec 7.6 #30

Hint: You may graph the solution using Mathematica.

11. Sec 7.6 #34

12. Free points!

Note: Mathematica commands are on the backside.

Mathematica Commands

See your HW 2 for the Mathematica commands. Below are some commands not in HW 2.

A piecewise defined function can be inputted using the `Piecewise` command;

`f[x_]:=Piecewise[{ {f1(x), a1 < x < b1}, {f2(x), a2 < x < b2}, ... }]`
is the function $f(x) = \begin{cases} f_1(x), & a_1 < x < b_1 \\ f_2(x), & a_2 < x < b_2 \\ \vdots \end{cases}$.

The Mathematica notation for the unit step function $u(t - a)$ is `UnitStep[t-a]`.

The Mathematica notation for the Dirac delta function $\delta(t - a)$ is `DiracDelta[t-a]`.

The Mathematica notation for the Laplace transform is `LaplaceTransform[f(t), t, s]` where $f(t)$ is the function and s is the independent variable of the Laplace transform $F(s)$.

The Mathematica notation for the inverse Laplace transform is `InverseLaplaceTransform[F(s), s, t]` where $F(s)$ is the Laplace transform and t is the independent variable of the inverse $f(t)$.