

Math 3280  
Review Problems for Exam I

This is not an exhaustive list of all possible types of problems. See the class notes, the textbook and homework for additional problems.

1. Solve the IVP  $x' + 2x = t e^{-2t}$ ,  $x(1) = 0$ .
2. Find the explicit solution of the IVP  $\frac{dx}{dt} = \frac{3t^2 + 4t + 2}{2(x-1)}$ ,  $x(0) = 2$ .
3. Find the implicit solution of  $\frac{dx}{dt} = \frac{x \cos t + 2t e^x}{1 - \sin t - t^2 e^x}$ .
4. Find the explicit solution  $x = x(t)$  of  $(t^2 + 3tx + x^2) dt - t^2 dx = 0$ . Assume  $t > 0$ .
5. Consider  $x'(t) = x - \sqrt[3]{4x}$ . Find its equilibrium solutions. Classify the equilibrium solutions and draw its phase line.
6. Consider  $x'(t) = x^2(1 - x^2)$ . Find its critical points and classify them using the linearization theorem. Draw its phase line.
7. Consider  $x'(t) = x^4 - 2x^3 + x^2$ . Find its equilibrium points and classify them. Determine the intervals, (initial)  $x$ -values, on which the solution  $x(t)$  would be increasing or decreasing. Determine the intervals,  $x$ -values, on which the solution  $x(t)$  would be concave up or down. Use these information to graph several representative flows (solutions) in the  $tx$ -plane.
8. Solve the IVP  $6x''(t) = 5x'(t) - x(t)$ ,  $x(0) = 4$ ,  $x'(0) = 0$  by first converting it to a first order planar system.
9. Consider the nonlinear system

$$\begin{aligned}\frac{dx}{dt} &= xy + 1 \\ \frac{dy}{dt} &= x^2 + 2\end{aligned}$$

Find all places in the  $xy$ -plane that its solution trajectories have (a) Horizontal tangent lines, (b) Vertical tangent lines, or (c) Tangent lines with slope  $m = 2$ , in any.

10. Solve

$$\begin{aligned}\frac{dx}{dt} &= -xy + e^{t^2} \\ \frac{dy}{dt} &= -\frac{y}{t}\end{aligned} \quad (x(1), y(1)) = \left(\frac{e}{2}, 1\right)$$

Assume  $t > 0$ .

(Partial) Answers to the Review Problems for Exam I Continued

17.  $\lambda = \lambda_1 = \lambda_2 = -1$ ,  $V_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $V_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;  $X(t) = c e^{-t} V$  for any vector  $V$ . All solutions are on lines through origin. The critical point  $(0, 0)$  is a sink.
18.  $\lambda = i$ ,  $V = \begin{pmatrix} 3 - i \\ 5 \end{pmatrix}$ ;  $X(t) = c_1 \begin{pmatrix} 3 \cos t + \sin t \\ 5 \cos t \end{pmatrix} + c_2 \begin{pmatrix} 3 \sin t - \cos t \\ 5 \sin t \end{pmatrix}$ . The critical point  $(0, 0)$  is a center. Solution curves have clockwise rotation.
19.  $\lambda = 1 + 2i$ ,  $V = \begin{pmatrix} -i \\ 1 \end{pmatrix}$ ;  $X(t) = c_1 \begin{pmatrix} e^t \sin 2t \\ e^t \cos 2t \end{pmatrix} + c_2 \begin{pmatrix} -e^t \cos 2t \\ e^t \sin 2t \end{pmatrix}$ . The critical point  $(0, 0)$  is a spiral source. Solution curves have clockwise rotation.
20.  $\lambda_1 = -2$ ,  $V_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ ;  $\lambda_2 = -1$ ,  $V_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ;  $X(t) = c_1 e^{-2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + c_2 e^{-t} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ . The lines  $y = x$  and  $y = 2x$  are the straight-line solutions (stable manifolds); The critical point  $(0, 0)$  is a sink.

11. Consider the nonlinear system

$$\begin{aligned}\frac{dx}{dt} &= 3y^2(x^2 + 1) \\ \frac{dy}{dt} &= 2x\end{aligned}$$

Solve it to find  $y$  as a function of  $x$  by considering  $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$ .

12. Find the solution of  $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$  with initial values  $x(0) = 2$  and  $y(0) = 1$

In the following problems, consider the system  $X'(t) = AX$ , with the given matrix  $A$ , and do the following.

- (a) Find its general solution.
- (b) Find its critical point(s) and classify them, if possible.
- (c) Discuss information useful for drawing its representative trajectories, for example,
  - Straight line (invariant) solutions.
  - Slopes of tangent lines to the solution curves.
  - Direction field or the direction of the increasing  $t$  value of the solution curves.
  - Converting the solution to rectangular coordinates.
- (d) Draw several representative trajectories.

Note: For these review problems, you may use Mathematica to find the eigenvalues and the corresponding eigenvectors.

$$13. A = \begin{pmatrix} 1 & -1 \\ -2 & 0 \end{pmatrix} \quad 14. A = \begin{pmatrix} 3 & -1 \\ 2 & 0 \end{pmatrix} \quad 15. A = \begin{pmatrix} 3 & -1 \\ 4 & -1 \end{pmatrix} \quad 16. A = \begin{pmatrix} 1 & -1 \\ 2 & -2 \end{pmatrix}$$

$$17. A = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \quad 18. A = \begin{pmatrix} -3 & 2 \\ -5 & 3 \end{pmatrix} \quad 19. A = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix} \quad 20. A = \begin{pmatrix} -3 & 1 \\ -2 & 0 \end{pmatrix}$$

Math 3280

(Partial) Answers to the Review Problems for Exam I

1.  $x(t) = \frac{1}{2}(t^2 - 1)e^{-2t}$
2.  $x(t) = 1 + \sqrt{t^3 + 2t^2 + 2t + 1}$
3.  $x - x \sin t - t^2 e^x = c$
4.  $x(t) = -\frac{(1+c)t + t \ln t}{c + \ln t}$
5.  $x = -2$ , unstable;  $x = 0$ , stable;  $x = 2$ , unstable
6.  $x = -1$ , unstable;  $x = 0$ , semistable;  $x = 1$ , stable
7.  $x = 0$  and  $x = 1$ , both semistable;  $x$  is increasing for  $x < 0$ ,  $0 < x < 1$ , and  $x > 1$ ;  $x$  is concave downward for  $x < 0$  and  $\frac{1}{2} < x < 1$ ;  $x$  is concave upward for  $0 < x < \frac{1}{2}$  and  $x > 1$
8.  $x = 12e^{\frac{1}{3}t} - 8e^{\frac{1}{2}t}$
9. No horizontal tangent lines; Vertical tangent lines on the curve  $y = -\frac{1}{x}$ ; Tangent lines with slope  $m = 2$  on the lines  $x = 0$  and  $y = \frac{x}{2}$
10.  $x(t) = \frac{e^{t^2}}{2t}$ ,  $y(t) = \frac{1}{t}$
11.  $y = -\sqrt[3]{c + \ln(x^2 + 1)}$
12.  $x(t) = (t + 2)e^{2t}$ ,  $y(t) = e^{2t}$
13.  $\lambda_1 = -1$ ,  $V_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ;  $\lambda_2 = 2$ ,  $V_2 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ ;  $X(t) = c_1 e^{-t} \begin{pmatrix} 1 \\ 2 \end{pmatrix} + c_2 e^{2t} \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ . The lines  $y = 2x$  and  $y = -x$  are the straight-line solutions (stable and unstable manifolds); The critical point  $(0, 0)$  is a saddle point.
14.  $\lambda_1 = 1$ ,  $V_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ;  $\lambda_2 = 2$ ,  $V_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ ;  $X(t) = c_1 e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} + c_2 e^{2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ . The lines  $y = 2x$  and  $y = x$  are the straight-line solutions (unstable manifolds); The critical point  $(0, 0)$  is a source.
15.  $\lambda = \lambda_1 = \lambda_2 = 1$ ,  $V = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ;  $X(t) = c_1 e^t \begin{pmatrix} 1 \\ 2 \end{pmatrix} + c_2 e^t \left( t \begin{pmatrix} 1 \\ 2 \end{pmatrix} + \begin{pmatrix} 0 \\ -1 \end{pmatrix} \right)$ ; The lines  $y = 2x$  is the straight-line solution (unstable manifolds); The critical point  $(0, 0)$  is a source; Solutions curves try to spiral about the critical point, but the unstable manifold gets in the way.
16.  $\lambda_1 = -1$ ,  $V_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ;  $\lambda_2 = 0$ ,  $V_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ ;  $X(t) = c_1 e^{-t} \begin{pmatrix} 1 \\ 2 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ . All points on the line  $y = x$  are critical points. All other solutions are lines with slope  $m = 2$  that move toward the line  $y = x$ .