

Challenges Faced by Homeless Sexual Minorities: Comparison of Gay, Lesbian, Bisexual, and Transgender Homeless Adolescents With Their Heterosexual Counterparts

Bryan N. Cochran, MS, Angela J. Stewart, BA, Joshua A. Ginzler, PhD, and Ana Mari Cauce, PhD

Homeless youths represent a diverse population that reaches the streets for a variety of reasons and whose numbers have grown in recent decades. Gay, lesbian, bisexual, and transgender (GLBT) homeless youths face the obstacle of survival on the streets as well as the stigma of sexual minority group membership.

It is difficult to estimate the proportion of GLBT youths in the street population.¹⁻³ The National Network of Runaway and Youth Services⁴ estimates that about 6% of homeless adolescents are gay or lesbian. However, the few studies assessing sexual orientations of homeless adolescents have revealed rates ranging from 11% to 35%.^{5,6}

Among adolescents in general, GLBT youths are more vulnerable to health and psychological problems than are heterosexual youths. Many are victims of parental physical abuse, are substance abusers, and have both mental and general physical health problems.⁷⁻¹⁰ These problems may be amplified for GLBT youths who become homeless.

Homeless youths are vulnerable to victimization, including robbery, rape, and assault.¹¹⁻¹⁵ Also, homelessness often leads to initiation or escalation of substance use.¹⁶⁻²⁴ High rates of externalizing and internalizing problems, including psychosis, have been found among this population.²⁵⁻³² Moreover, high rates of risky sexual behavior, including prostitution and survival sex (sex in exchange for money, drugs, or shelter), place these young people at risk for victimization and sexually transmitted diseases.³³⁻³⁶

The current study was the first of its kind to examine psychosocial outcomes for GLBT homeless youths. The objectives of the study were to identify risks faced by GLBT homeless youths and to determine whether these risks transcend those of their heterosexual counterparts. Our hypothesis was that GLBT home-

Objectives. The goal of this study was to identify differences between gay, lesbian, bisexual, and transgender (GLBT) homeless youths and their heterosexual counterparts in terms of physical and mental health difficulties.

Methods. A sample of 84 GLBT adolescents was matched in regard to age and self-reported gender with 84 heterosexual adolescents. The 2 samples were compared on a variety of psychosocial variables.

Results. GLBT adolescents left home more frequently, were victimized more often, used highly addictive substances more frequently, had higher rates of psychopathology, and had more sexual partners than heterosexual adolescents.

Conclusions. Homeless youths who identify themselves as members of sexual minority groups are at increased risk for negative outcomes. Recommendations for treatment programs and implications for public health are discussed. (*Am J Public Health.* 2002; 92:773-777)

less youths would have more negative indicators of psychological and physical well-being than do heterosexual homeless adolescents.

METHOD

Participants

The sample consisted of 375 adolescents aged 13 to 21 years (mean: 17.14 years); data were collected between 1995 and 1998. Youths were recruited for the Seattle Homeless Adolescent Research and Education project at street locations or social service agencies in the Seattle metropolitan area. Youths were eligible to participate if they spoke English, had not lived in the residence of a primary caretaker for at least 1 week, and had no stable home in which to live.

Trained outreach workers informed youths of their right to refuse participation, skip individual questions, or stop participation at any time. Informed consent was obtained from all participants, and youths were offered \$25 for taking part. The overall response rate was 95%. The sampling methods were described in full in an earlier report.³⁷

The majority of study participants identified themselves as White (52.5%). Other par-

ticipants self-identified as American Indian or Alaska Native (18.9%), African American (17.6%), Hispanic/Latino (7.2%), or Asian/Pacific Islander (2.7%). About half of the sample was female (45.1%).

Self-reports of sexual orientation (heterosexual, gay, lesbian, bisexual, transsexual, or transgender) were used to classify the majority of participants. In a few cases, youths did not select one of the choices ($n=9$) or selected multiple choices ($n=12$). Questions about same- and opposite-sex attraction were used to classify these individuals as members of sexual minorities ($n=6$) or as heterosexuals ($n=15$). Individuals who both rated their same-sex attraction higher than the midpoint on a 7-point Likert scale and reported same-sex sexual behavior were classified as sexual minorities.

The majority of the 84 sexual minority youths identified themselves as bisexual ($n=71$). Only 4 female and 8 male youths self-identified as exclusively lesbian or gay, and 1 youth self-identified as transgender. GLBT youths were matched in terms of age and gender with an equal number of self-identified heterosexual participants; thus, the study included a total of 168 participants.

Measures

Private, face-to-face structured interviews (1.5 to 2 hours in duration) were conducted with each youth. Participants were asked about their reasons for leaving home, street victimization, alcohol and drug use, and sexual behaviors. In all, 21 reasons for leaving home were provided, and youths were asked to select any reason that influenced their decision to leave. Youths were also asked whether they had encountered any physical or sexual victimization after they had left home, and if so, how often it had occurred (e.g., how often they had been assaulted or forced to have sex).

In the substance use section, participants were asked how often they had used all major forms of drugs and alcohol in the previous 6 months (ratings were made on a 7-point scale ranging from *not at all* [0] to *every day* [6]). Sexual behavior questions focused on areas such as number and gender of sexual partners and frequency of safe-sex practices. Depressive symptoms were measured with the Center for Epidemiologic Studies Depression Scale (CES-D),^{38,39} and other behavior problems were assessed with Achenbach's Youth Self-Report (YSR).⁴⁰ A full description of measures is available from the authors.

Data Analyses

Given our primary goal of determining how GLBT youths differed from heterosexual youths in several domains, we had no a priori multivariate predictions about the relationships among variables. Therefore, we conducted primarily *t* tests (for continuous variables) and, in some cases, χ^2 tests (for dichotomous variables).

RESULTS

Pathways to Homelessness

By and large, GLBT youths left home for reasons similar to those of their heterosexual counterparts. However, GLBT youths left home more often than did heterosexual youths (means of 12.38 times and 6.69 times, respectively; $t_{160} = 1.91, P = .058$). The most common reasons reported by youths for leaving home were family conflict (59.9%), desire for freedom (51.5%),

TABLE 1—Physical and Sexual Victimization of Sexual Minority and Heterosexual Homeless Adolescents: Seattle, Wash, 1995–1998

Measure	Sexual Minorities, Mean (SD)	Heterosexuals, Mean (SD)	<i>t</i> (df)	<i>P</i>
Physical victimization				
Since homeless	1.88 (0.70)	1.66 (0.72)	1.98 (165)	.05
Male	2.10 (0.72)	2.00 (0.77)	0.58 (73)	.56
Female	1.70 (0.64)	1.39 (0.54)	2.49 (90)	.02
During past 3 months	1.45 (0.50)	1.25 (0.44)	2.67 (166)	.008
Male	1.67 (0.58)	1.39 (0.55)	2.21 (74)	.03
Female	1.26 (0.32)	1.14 (0.29)	1.87 (90)	.07
Sexual victimization				
Since homeless	1.52 (0.75)	1.27 (0.52)	2.46 (161)	.02
Male	1.36 (0.68)	1.05 (0.16)	2.65 (71)	.01
Female	1.65 (0.78)	1.46 (0.63)	1.31 (88)	.19
During past 3 months	1.45 (0.82)	1.24 (0.56)	1.80 (138)	.07
Male	1.50 (0.99)	1.09 (0.37)	2.25 (65)	.03
Female	1.40 (0.63)	1.37 (0.66)	0.21 (71)	.83
No. of perpetrators	8.61 (24.06)	1.24 (3.76)	2.69 (156)	.009
Male	6.74 (22.95)	0.17 (0.45)	1.70 (69)	.01
Female	10.09 (25.07)	2.14 (4.93)	2.06 (85)	.05

Note. For all measures except number of perpetrators, possible responses ranged from 1 (never) to 4 (many times).

and difficulties with a family member (48.5%). GLBT youths were more likely to leave as a result of physical abuse in the home ($\chi^2_1 = 3.6, P = .044$), and there was a trend toward more GLBT youths leaving as a result of alcohol use in the home ($\chi^2_1 = 3.2, P = .055$). Twelve (14.3%) GLBT youths indicated that they had left home because of conflicts with their parents over their sexual orientation.

Victimization


GLBT youths experienced higher levels of physical victimization than their heterosexual counterparts (Table 1). When analyses were conducted separately by gender, this effect was significant for male youths during the preceding 3 months and for female youths since the onset of their homelessness. Also, GLBT youths had more often been sexually victimized since the time they had first become homeless. When examined by gender, this effect was found among male youths but not among female youths. In addition, GLBT youths reported an average of 7.4 more perpetrators of sexual victimization than did heterosexual youths.

Substance Abuse

For each substance category except marijuana, sexual minority youths had used the substance more frequently in the previous 6 months than had heterosexuals (Figure 1). Differences were significant for 3 types of drugs: cocaine or crack (respective means of 0.81 and 0.33 on the 7-point scale; $t_{166} = 2.75, P = .007$), crack or crack mixed with amphetamines (means of 1.02 and 0.46, respectively; $t_{166} = 2.43, P = .016$), and speed or crystal methamphetamines (means of 1.56 and 0.74; $t_{166} = 2.88, P = .005$). Sexual minority youths also used more types of drugs than did heterosexuals (means of 7.88 and 6.51, respectively; $t_{166} = 2.39, P = .018$).


Mental Health

There was a trend for GLBT youths to report higher levels of depressive symptoms (as measured with the CES-D) than did heterosexual youths ($t_{147} = 1.43, P = .09$). Also, YSR subscale scores indicated that GLBT youths had higher rates of psychopathology (Figure 2). Independent-samples *t* tests revealed significant differences in the following domains: withdrawn behavior ($t_{142} = 2.28, P = .024$), somatic



Note. * $P < .05$; ** $P < .01$.

FIGURE 1—Average use of substances in the past 6 months as a function of sexual orientation: Seattle, Wash, 1995–1998



Note. * $P < .05$; ** $P < .01$.

FIGURE 2—Profiles of Youth Self-Report (YSR) t scales, by sexual orientation: Seattle, Wash, 1995–1998

complaints ($t_{141}=2.07, P=.041$), social problems ($t_{150}=2.03, P=.044$), delinquency ($t_{142}=2.07, P=.040$), aggression ($t_{149}=2.45, P=.016$), internalizing behavior ($t_{146}=2.38, P=.018$), and externalizing behavior ($t_{148}=2.71, P=.007$). The total YSR mean difference was also significant; overall, GLBT adolescents re-

ported higher levels of symptomatology ($t_{149}=3.24, P=.001$).

Sexual Behavior

Most adolescents (94%) reported engaging in voluntary sex at least once; however, GLBT youths reported a higher number of

lifetime sexual partners than did heterosexual youths (means of 24.19 and 12.49, respectively; $t_{143}=2.84, P=.005$). GLBT youths were significantly younger than heterosexuals in regard to age at first voluntary intercourse (means of 13.00 and 13.83 years, respectively; $t_{154}=2.11, P=.037$). Also, GLBT youths reported high rates of unprotected intercourse, endorsing a mean of 2.91 for this item (approximately “half of the time”), in comparison with a mean of 2.51 among heterosexual adolescents ($t_{115}=1.08, NS$). Although this difference was not statistically significant, more than twice as many GLBT youths as heterosexual youths reported that they neglected to use protection during sex “all of the time.”

DISCUSSION

The present results indicate negative outcomes in multiple domains for GLBT homeless adolescents. These outcomes include more-frequent departures from home, greater vulnerability to physical and sexual victimization, higher rates of addictive substance use, more psychopathology, and riskier sexual behavior in comparison with homeless heterosexual adolescents.

The coping model used by MacLean et al.²¹ to explain substance use among homeless adolescents may be extended to these findings. GLBT homeless adolescents experience not only the vulnerabilities, daily difficulties, and survival challenges of living on the street but also the discrimination faced by GLBT youth in general. In coping with these stressors, they may use more substances and use them more frequently than do heterosexual youths. Other psychological problems may be direct effects of this discrimination or may be intensified by coping-focused substance abuse.

GLBT adolescents face great challenges as they work to come to terms with their sexual orientation. Those living in homes forge this identity within the context of their family and school or peer networks, facing risks of isolation, rejection, and sometimes victimization by others. Their homeless counterparts, however, frequently have no family members available, no school environment to support them, and transient or insufficient peer networks.

Primary interventions directed toward these youths should focus on preventing initial and recurrent episodes of homelessness by providing therapeutic services to assist families in dealing with adolescents' sexual identity, thus improving the home environment and reducing the likelihood that GLBT adolescents will leave. Secondary interventions that could benefit GLBT homeless adolescents would involve recognition of the particular difficulties faced by this group and provision of services sensitive to issues of sexual orientation.

Community public health programs must recognize the increased risks faced by homeless GLBT youths and the need for assisting this population. First, it is important to ask adolescents about their sexual orientation, both to identify their need for services and to indicate that sexual orientation is an acceptable topic of discussion. Second, intensified services similar to those described by Tenner et al.⁶ could be tailored to homeless GLBT youths to reduce risks of HIV infection. Third, programs should acknowledge and address the contribution of homophobia to the etiology and maintenance of substance abuse problems. Finally, programs that encourage acceptance of sexual minorities among street youths and in shelters may reduce the risk of GLBT adolescents being victimized in these locales. Further studies may increase our understanding of this often-overlooked population.

Our study involved some limitations that should be noted. All of the data were obtained through self-reports, and most domains were assessed via single measurements. Our sexual minority group was largely bisexual, and results may not generalize to homeless adolescents who self-identify as exclusively lesbian, gay, or transgender. However, sexual orientation is a dynamic variable subject to developmental changes,³ and future studies may elucidate whether and how outcomes change as homeless adolescents redefine their sexual identities. Finally, our analyses were limited to bivariate tests. Given our interest in exploring differences between sexual minority and heterosexual homeless adolescents in a number of different domains, our approach was appropriate; in the future, however, we will use multivariate modeling techniques in

an attempt to improve our understanding of the challenges faced by GLBT youths in the context of homelessness. ■

About the Authors

Bryan N. Cochran and Angela J. Stewart are with the Department of Psychology, University of Washington, Seattle. Joshua A. Ginzler is with the Alcohol and Drug Abuse Institute, Seattle. Ana Mari Cauce is with the Departments of Psychology and American Ethnic Studies, University of Washington.

Requests for reprints should be sent to Bryan N. Cochran, MS, Box 351525, University of Washington, Seattle, WA 98195 (e-mail: cochranb@u.washington.edu). This article was accepted December 31, 2001.

Contributors

B.N. Cochran conceptualized the project, conducted data analyses, and contributed to the literature review and the writing of the manuscript. A.J. Stewart contributed to the study design, to the literature review and data analysis, and to the writing and review of the manuscript. J.A. Ginzler assisted with the substance abuse outcome data analysis and the literature review and linked findings to the risk amplification model. A.M. Cauce contributed to study design and implementation and to the writing and review of the manuscript.

Acknowledgments

This study was supported by a research grant (AA10253-03) from the National Institute on Alcohol Abuse and Alcoholism.

We thank Brynn Malone for her contributions to the editing of the manuscript and Kevin Wruck and Yvette Lohr for their conscientious data collection efforts.

References

- Sell RL. Defining and measuring sexual orientation: a review. *Arch Sex Behav*. 1997;26:643-658.
- Rotheram-Borus MJ, Koopman C, Ehrhardt AA. Homeless youths and HIV infection. *Am Psychol*. 1991;46:1188-1197.
- Diamond LM. Sexual identity, attractions, and behavior among young sexual-minority women over a 2-year period. *Dev Psychol*. 2000;36:241-250.
- To Whom Do They Belong? A Profile of America's Runaway and Homeless Youth and the Programs That Help Them*. Washington, DC: National Network of Runaway and Youth Services; 1985.
- Kruks G. Gay and lesbian homeless/street youth: special issues and concerns. *J Adolesc Health*. 1991;12:515-518.
- Tenner AD, Trevithick LA, Wagner V, Burch R. Seattle YouthCare's prevention, intervention and education program: a model of care for HIV-positive, homeless, and at-risk youth. *J Adolesc Health*. 1998;23:96-106.
- Savin-Williams RC, Cohen KM. Psychosocial outcomes of verbal and physical abuse among lesbian, gay, and bisexual youths. In: Savin-Williams RC, Cohen KM, eds. *The Lives of Lesbians, Gays, and Bisexuals: Children to Adults*. Fort Worth, Tex: Harcourt Brace College Publishers; 1996:181-200.
- Rosario M, Hunter J, Gwadz M. Exploration of substance use among lesbian, gay, and bisexual youth: prevalence and correlates. *J Adolesc Res*. 1997;12:454-476.
- Fergusson DM, Horwood LJ, Beautrais AL. Is sexual orientation related to mental health problems and suicidality in young people? *Arch Gen Psychiatry*. 1999;56:876-880.
- Lock J, Steiner H. Gay, lesbian, and bisexual youth risks for emotional, physical, and social problems: results from a community-based survey. *J Am Acad Child Adolesc Psychiatry*. 1999;38:297-304.
- Baron SW. Risky lifestyles and the link between offending and victimization. *Stud Crime Crime Prev*. 1997;6:53-71.
- Hoyt DR, Ryan KD, Cauce AM. Personal victimization in a high-risk environment: homeless and runaway adolescents. *J Res Crime Delinquency*. 1999;36:371-392.
- Kipke MD, Simon TR, Montgomery SB, Unger JB, Iversen EF. Homeless youth and their exposure to and involvement in violence while living on the streets. *J Adolesc Health*. 1997;20:360-367.
- Whitbeck LB, Hoyt DR, Ackley KA. Abusive family backgrounds and later victimization among runaway and homeless adolescents. *J Res Adolesc*. 1997;7:375-392.
- Whitbeck LB, Simons RL. A comparison of adaptive strategies and patterns of victimization among homeless adolescents and adults. *Violence Vict*. 1993;8:135-152.
- Fors SW, Rojek DG. A comparison of drug involvement between runaways and school youths. *J Drug Educ*. 1991;21:13-25.
- Forst ML. A substance use profile of delinquent and homeless youths. *Int J Psychiatry Med*. 1994;24:219-231.
- Greene JM, Ennett ST, Ringwalt CL. Substance abuse among runaway and homeless youth in three national samples. *Am J Public Health*. 1997;87:229-235.
- Kipke MD, Montgomery S, MacKenzie RG. Substance use among youth seen at a community-based health clinic. *J Adolesc Health*. 1993;14:289-294.
- Koopman C, Rosario M, Rotheram-Borus MJ. Alcohol and drug use and sexual behaviors placing runaways at risk for HIV infection. *Addict Behav*. 1994;19:95-103.
- MacLean MG, Paradise MJ, Cauce AM. Substance use and psychological adjustment in homeless adolescents: a test of three models. *Am J Community Psychol*. 1999;27:405-427.
- Robertson MJ, Koegel P, Ferguson L. Alcohol use and abuse among homeless adolescents in Hollywood. *Contemp Drug Problems*. 1989;16:415-452.
- Smart RG, Adlaf EM. Substance use and problems among Toronto street youth. *Br J Addict*. 1991;86:999-1010.
- Unger JB, Kipke MD, Simon TR, Montgomery SB, Johnson CJ. Homeless youths and young adults in Los Angeles: prevalence of mental health problems and the relationship between mental health and substance abuse disorders. *Am J Community Psychol*. 1997;25:371-394.
- Cauce AM, Paradise M, Ginzler JA, et al. The

characteristics and mental health of homeless adolescents: age and gender differences. *J Emotional Behav Disord.* 2000;8:230-239.

26. Feitel B, Margetson N, Chamas R, Lipman C. Psychosocial background and behavioral and emotional disorders of homeless and runaway youth. *Hosp Community Psychiatry.* 1992;43:155-159.

27. Fosberg LB, Dennis DL. *Practical Lessons: The 1998 National Symposium on Homelessness Research.* Washington, DC: US Dept of Housing and Urban Development; 1999.

28. Greene JM, Ringwalt CL. Youth and familial substance use's association with suicide attempts among runaway and homeless youth. *Subst Use Misuse.* 1996;31:1041-1058.

29. McCaskill PA, Toro PA, Wolfe SM. Homeless and matched housed adolescents: a comparative study of psychopathology. *J Clin Child Psychol.* 1998;27:306-319.

30. Mundy P, Robertson M, Robertson J, Greenblatt M. The prevalence of psychotic symptoms in homeless adolescents. *J Am Acad Child Adolesc Psychiatry.* 1990;29:724-731.

31. Rotheram-Borus MJ. Suicidal behavior and risk factors among runaway youths. *Am J Psychiatry.* 1993;150:103-107.

32. Yoder KA, Hoyt DR, Whitbeck LB. Suicidal behavior among homeless and runaway adolescents. *J Youth Adolesc.* 1998;27:753-771.

33. Greenblatt M, Robertson MJ. Life-styles, adaptive strategies, and sexual behaviors of homeless adolescents. *Hosp Community Psychiatry.* 1993;44:1177-1180.

34. Remafedi GR. Adolescent homosexuality: psychosocial and medical implications. *Pediatrics.* 1987;79:331-337.

35. Pennbridge JN, Freese TE, MacKenzie RG. High-risk behaviors among male street youth in Hollywood, California. *AIDS Educ Prev.* Fall 1992(suppl):24-33.

36. Yates GL, MacKenzie RG, Pennbridge J, Swofford A. A risk profile comparison of homeless youth involved in prostitution and homeless youth not involved. *J Adolesc Health.* 1991;12:545-548.

37. Tyler KA, Hoyt DR, Whitbeck LB, Cauce AM. The impact of childhood sexual abuse on later sexual victimization among runaway youth. *J Res Adolesc.* 2001;11:151-176.

38. Radloff LS. The CES-D Scale: a self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977;1:385-401.

39. Radloff LS. The use of the Center for Epidemiologic Studies Depression Scale in adolescents and young adults. *J Youth Adolesc.* 1991;20:149-166.

40. Achenbach TM. *Manual for the Youth Self-Report and 1991 Profile.* Burlington, Vt: University of Vermont; 1991.

Community-Oriented Primary Care: Health Care for the 21st Century



2nd Edition

ISBN 0-87553-236-5

1998 ■ 228 pages

Softcover

\$27.00 APHA Members

\$39.00 Nonmembers

plus shipping and handling

Community-Oriented Primary Care: Health Care for the 21st Century

Edited by Robert Rhyne, MD, Richard Bogue, PhD,
Gary Kukulka, PhD, and Hugh Fulmer, MD

This book will give insight into:

- How medicine, health systems, community leaders, and social services can be supportive as America's public health practice continues to be restructured and redefined
- New models of community-oriented primary care
- Methods and interventions on population-derived health needs
- Health promotion and disease prevention as part of the overall reorganization of health services
- Understanding how community-oriented primary care can complement managed care and community benefit programs

This book teaches skills and techniques for implementing a community-oriented primary care process and topics not normally taught in health professional education.



American Public Health Association

Publication Sales

Web: www.apha.org

E-mail: APHA@TASCO1.com

Tel: (301) 893-1894

FAX: (301) 843-0159

COPC03J5

Copyright of American Journal of Public Health is the property of American Public Health Association and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.